

Rec'd PCTO

532, 426  
22 APR 2005

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
6 May 2004 (06.05.2004)

PCT

(10) International Publication Number  
**WO 2004/037277 A2**

(51) International Patent Classification<sup>7</sup>: **A61K 38/00**

(21) International Application Number:  
PCT/CA2003/001635

(22) International Filing Date: 24 October 2003 (24.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/420,677 24 October 2002 (24.10.2002) US

(71) Applicant (for all designated States except US): **MCGILL UNIVERSITY** [CA/CA]; 845 Sherbrooke Street West, Montréal, Québec H3A 2T5 (CA).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **ROSENBERG, Lawrence** [CA/CA]; 6507 Fern Road, Côte St-Luc, Québec H4V 1E4 (CA).

(74) Agent: **OGILVY RENAULT**; Suite 1600, 1981 McGill College Avenue, Montreal, Québec H3A 2Y3 (CA).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND COMPOSITION FOR REVERSAL OF DIABETES AND USES THEREOF

(57) Abstract: The present invention relates to a method to stimulate reversal of a diabetic state in a patient; a method to prevent autoimmune destruction of new insulin-producing cells (pancreatic beta-cells) in a patient; a method to promote survival of the newly regenerated insulin-producing cells (pancreatic beta-cells); and an in vivo method for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells.

WO 2004/037277 A2

## METHODS AND COMPOSITION FOR REVERSAL OF DIABETES AND USES THEREOF

### BACKGROUND OF THE INVENTION

#### 5 (a) Field of the Invention

This invention relates to a method to stimulate reversal of a diabetic state in a patient; a method to prevent autoimmune destruction of new insulin-producing cells (pancreatic beta-cells) in a patient; a method to promote survival of the newly regenerated insulin-producing cells  
10 (pancreatic beta-cells); and an *in vivo* method for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells; pharmaceutical compositions and uses thereof.

#### (b) Description of Prior Art

#### 15 **Diabetes**

Diabetes mellitus has been classified as type I, or insulin-dependent diabetes mellitus (IDDM) and type II, or non-insulin-dependent diabetes mellitus (NIDDM). NIDDM patients have been subdivided further into (a) nonobese (possibly IDDM in evolution), (b) obese, and (c) maturity  
20 onset (in young patients). Among the population with diabetes mellitus, about 20% suffer from IDDM. Diabetes develops either when a diminished insulin output occurs or when a diminished sensitivity to insulin cannot be compensated for by an augmented capacity for insulin secretion. In patients with IDDM, a decrease in insulin secretion is the principal factor in  
25 the pathogenesis, whereas in patients with NIDDM, a decrease in insulin sensitivity is the primary factor. The mainstay of diabetes treatment, especially for type I disease, has been the administration of exogenous insulin.

#### 30 **Rationale for more physiologic therapies**

Tight glucose control appears to be the key to the prevention of the secondary complications of diabetes. The results of the Diabetes Complications and Control Trial (DCCT), a multicenter randomized trial of  
35 1441 patients with insulin dependent diabetes, indicated that the onset and progression of diabetic retinopathy, nephropathy, and neuropathy could be

slowed by intensive insulin therapy (The Diabetes Control and  
Complication Trial Research Group, *N. Engl. J. Med.*, 1993; 29:977-986).  
Strict glucose control, however, was associated with a three-fold increase  
in incidence of severe hypoglycemia, including episodes of seizure and  
coma. As well, although glycosylated hemoglobin levels decreased in the  
5 treatment group, only 5% maintained an average level below 6.05%  
despite the enormous amount of effort and resources allocated to the  
support of patients on the intensive regime (The Diabetes Control and  
Complication Trial Research Group, *N. Engl. J. Med.*, 1993; 29:977-986).  
10 The results of the DCCT clearly indicated that intensive control of glucose  
can significantly reduce (but not completely protect against) the long-term  
microvascular complications of diabetes mellitus.

#### Other therapeutic options

15 The delivery of insulin in a physiologic manner has been an  
elusive goal since insulin was first purified by Banting, Best, McLeod and  
Collip. Even in a patient with tight glucose control, however, exogenous  
insulin has not been able to achieve the glucose metabolism of an  
endogenous insulin source that responds to moment-to-moment changes  
20 in glucose concentration and therefore protects against the development of  
microvascular complications over the long term.

A major goal of diabetes research, therefore, has been the  
development of new forms of treatment that endeavor to reproduce more  
closely the normal physiologic state. One such approach, a closed-loop  
25 insulin pump coupled to a glucose sensor, mimicking  $\beta$ -cell function in  
which the secretion of insulin is closely regulated, has not yet been  
successful. Only total endocrine replacement therapy in the form of a  
transplant has proven effective in the treatment of diabetes mellitus.  
Although transplants of insulin-producing tissue are a logical advance over  
30 subcutaneous insulin injections, it is still far from clear whether the risks of  
the intervention and of the associated long-term immunosuppressive  
treatment are lower those in diabetic patients under conventional  
treatment.

Despite the early evidence of the potential benefits of  
35 vascularized pancreas transplantation, it remains a complex surgical

intervention, requiring the long-term administration of chronic immunosuppression with its attendant side effects. Moreover, almost 50% of successfully transplanted patients exhibit impaired tolerance curves (Wright FH et al., *Arch. Surg.*, 1989; **124**:796-799; Landgraft R et al., *Diabetologia* 1991; **34** (suppl 1):S61; Morel P et al., *Transplantation* 1991; **51**:990-1000), raising questions about their protection against the long-term complications of chronic hyperglycemia.

The major complications of whole pancreas transplantation, as well as the requirement for long term immunosuppression, has limited its wider application and provided impetus for the development of islet transplantation. Theoretically, the transplantation of islets alone, while enabling tight glycemic control, has several potential advantages over whole pancreas transplantation. These include the following: (i) minimal surgical morbidity, with the infusion of islets directly into the liver via the portal vein; (ii) the possibility of simple re-transplantation for graft failures; (iii) the exclusion of complications associated with the exocrine pancreas; (iv) the possibility that islets are less immunogenic, eliminating the need for immunosuppression and enabling early transplantation into non-uremic diabetics; (v) the possibility of modifying islets in vitro prior to transplantation to reduce their immunogenicity; (vi) the ability to encapsulate islets in artificial membranes to isolate them from the host immune system; and (vii) the related possibility of using xenotransplantation of islets immunoisolated as part of a biohybrid system. Moreover, they permit the banking of the endocrine cryopreserved tissue and a careful and standardized quality control program before the implantation.

### **The problem of islet transplantation**

Adequate numbers of isogenetic islets transplanted into a reliable implantation site can only reverse the metabolic abnormalities in diabetic recipients in the short term. In those that were normoglycemic post-transplant, hyperglycemia recurred within 3-12 mo. (Orloff M, et. al., *Transplantation* 1988; **45**:307). The return of the diabetic state that occurs with time has been attributed either to the ectopic location of the islets, to a disruption of the enteroinsular axis, or to the transplantation of an

inadequate islet cell mass (Bretzel RG, et al. In: Bretzel RG, (ed) Diabetes mellitus (Berlin: Springer, 1990) p.229).

5       Studies of the long term natural history of the islet transplant, that  
examine parameters other than graft function, are few in number. Only one  
report was found in which an attempt was specifically made to study graft  
morphology (Alejandro R, et. al., *J Clin Invest* 1986; **78**: 1339). In that  
study, purified islets were transplanted into the canine liver via the portal  
vein. During prolonged follow-up, delayed failures of graft function  
occurred. Unfortunately, the graft was only examined at the end of the  
10 study, and not over time as function declined. Delayed graft failures have  
also been confirmed by other investigators for dogs (Warnock GL et. al.,  
*Can. J. Surg.*, 1988; **31**: 421 and primates (Sutton R, et. al., *Transplant  
Proc.*, 1987; **19**: 3525). Most failures are presumed to be the result of  
rejection despite appropriate immunosuppression.

15       Because of these failures, there is currently much enthusiasm for  
the immunoisolation of islets, which could eliminate the need for  
immunosuppression. The reasons are compelling. Immunosuppression is  
harmful to the recipient, and may impair islet function and possibly cell  
survival (Metrakos P, et al., *J. Surg. Res.*, 1993; **54**: 375). Unfortunately,  
20 micro-encapsulated islets injected into the peritoneal cavity of the dog fail  
within 6 months (Soon-Shiong P, et. al., *Transplantation* 1992; **54**: 769),  
and islets placed into a vascularized biohybrid pancreas also fail, but at  
about one year. In each instance, however, histological evaluation of the  
graft has indicated a substantial loss of islet mass in these devices (Lanza  
25 RP, et. al., *Diabetes* 1992; **41**: 1503). No reasons have been advanced for  
these changes. Therefore maintenance of an effective islet cell mass post-  
transplantation remains a significant problem.

30       In addition to this unresolved issue, is the ongoing problem of the  
lack of source tissue for transplantation. The number of human donors is  
insufficient to keep up with the potential number of recipients. Moreover,  
given the current state of the art of islet isolation, the number of islets that  
can be isolated from one pancreas is far from the number required to  
effectively reverse hyperglycemia in a human recipient.

In response, three competing technologies have been proposed and are under development. First, islet cryopreservation and islet banking. The techniques involved, though, are expensive and cumbersome, and do not easily lend themselves to widespread adoption. In addition, islet cell mass is also lost during the freeze-thaw cycle. Therefore this is a poor long-term solution to the problem of insufficient islet cell mass. Second, is the development of islet xenotransplantation. This idea has been coupled to islet encapsulation technology to produce a biohybrid implant that does not, at least in theory, require immunosuppression. There remain many problems to solve with this approach, not least of which, is that the problem of the maintenance of islet cell mass in the post-transplant still remains. Third, is the resort to human fetal tissue, which should have a great capacity to be expanded *ex vivo* and then transplanted. However, in addition to the problems of limited tissue availability, immunogenicity, there are complex ethical issues surrounding the use of such a tissue source that will not soon be resolved. However, there is an alternative that offers similar possibilities for near unlimited cell mass expansion.

An entirely novel approach, proposed by Rosenberg in 1995 (Rosenberg L et al., *Cell Transplantation*, 1995; 4:371-384), was the development of technology to control and modulate islet cell neogenesis and new islet formation, both *in vitro* and *in vivo*. The concept assumed that (a) the induction of islet cell differentiation was in fact controllable; (b) implied the persistence of a stem cell-like cell in the adult pancreas; and (c) that the signal(s) that would drive the whole process could be identified and manipulated.

In a series of *in vivo* studies, Rosenberg and co-workers established that these concepts were valid in principle, in the *in vivo* setting (Rosenberg L et al., *Diabetes*, 1988; 37:334-341; Rosenberg L et al., *Diabetologia*, 1996; 39:256-262), and that diabetes could be reversed.

The well known teachings of *in vitro* islet cell expansion from a non-fetal tissue source comes from Peck and co-workers (Corneliu JG et al., *Horm. Metab. Res.*, 1997; 29:271-277), who describe isolation of a pluripotent stem cell from the adult mouse pancreas that can be directed toward an insulin-producing cell. These findings have not been widely

accepted. First, the result has not proven to be reproducible. Second, the so-called pluripotential cells have never been adequately characterized with respect to phenotype. And third, the cells have certainly not been shown to be pluripotent.

5 More recently two other competing technologies have been proposed the use of engineered pancreatic  $\beta$ -cell lines (Efrat S, *Advanced Drug Delivery Reviews*, 1998; 33:45-52), and the use of pluripotent embryonal stem cells (Shamblott MJ et al., *Proc. Natl. Acad. Sci. USA*, 1998; 95:13726-13731). The former option, while attractive, is associated  
10 with significant problems. Not only must the engineered cell be able to produce insulin, but it must respond in a physiologic manner to the prevailing level of glucose- and the glucose sensing mechanism is far from being understood well enough to engineer it into a cell. Many proposed cell lines are also transformed lines, and therefore have a neoplastic potential.  
15 With respect to the latter option, having an embryonal stem cell in hand is appealing because of the theoretical possibility of being able to induce differentiation in any direction, including toward the pancreatic  $\beta$ -cell. However, the signals necessary to achieve this milestone remain unknown.

Islet neogenesis associated protein (INGAP) is a mediator of *in vivo* islet cell neogenesis from pancreatic duct epithelial cells in several species.  
20

It would be highly desirable to be provided with a method for the *in vivo* induction of re-growth of new insulin-producing cells leading to the formation of mature islets of Langerhans using INGAP peptide (the  
25 biologically active portion of the INGAP molecule), as a means of revering an established diabetic state.

Moreover, if such a diabetic was caused by pre-existing or ongoing autoimmunity, it would also be highly desirable to be provided with a method for the mitigation of such autoimmunity so that the  
30 aforementioned newly re-grown cells will not be subjected to ongoing or renewed destruction.

## SUMMARY OF THE INVENTION

One aim of the invention is to provide a method to stimulate reversal of a diabetic state in a patient.

Another aim of the invention is to provide a method to prevent  
5 autoimmune destruction of new insulin-producing cells (pancreatic beta-cells) in a patient.

Another aim of the invention is to provide a method to promote survival of the newly regenerated insulin-producing cells (pancreatic beta-cells).

10 Another aim of the invention is to provide an *in vivo* method for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of the new cells.

In accordance with the present invention there is provided a method to stimulate reversal of a diabetic state in a patient, which  
15 comprises *in vivo* inducing re-growth of new insulin-producing cells by administering a therapeutically effective amount of a pro-neogenesis factor to said patient, wherein formation of mature islets of Langerhans is indicative of a stimulated reversal of a diabetic state.

In accordance with the present invention there is provided a  
20 method to prevent autoimmune destruction of new insulin-producing cells (pancreatic beta-cells) in a patient, which comprises administering to said patient a therapeutically effective amount of at least one immunosuppressive agent in combination with an INGAP peptide.

In accordance with the present invention there is provided an *in*  
25 *vivo* method for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells, which comprises the steps of:

a) administering INGAP peptide to said patient in an amount  
30 sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms, whereby cells expand in number and develop a mature glucose-sensing mechanism in a regulated manner;



- b) concurrently administering to said patient at least one immunosuppressive agent in an amount sufficient to protect said islet cells from immune destruction; and
- c) concurrently administering a pro-survival factor to said patient during islet cell neogenesis and new islet formation.

In accordance with the present invention there is provided an *in vivo* method for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells, which comprises the steps of:

- a) administering INGAP peptide to said patient in an amount sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms, whereby cells expand in number and develop a mature glucose-sensing mechanism in a regulated manner;
- b) concurrently administering a pro-survival factor to said patient during islet cell neogenesis and new islet formation.

In accordance with the present invention there is provided a pharmaceutical composition for the preparation of a medicament to stimulate reversal of a diabetic state in a patient by *in vivo* inducing re-growth of new insulin-producing cells, which comprises a therapeutically effective amount of a pro-neogenesis factor in association with a pharmaceutically acceptable carrier.

In accordance with the present invention there is provided a pharmaceutical composition for the preparation of a medicament to prevent autoimmune destruction of new insulin-producing cells in a patient, which comprises a therapeutically effective amount of at least one immunosuppressive agent and an INGAP peptide factor in association with a pharmaceutically acceptable carrier.

In accordance with the present invention there is provided a pharmaceutical composition for the preparation of a medicament to promote survival of the newly regenerated insulin-producing cells, which comprises a therapeutically effective amount of a pro-neogenesis factor in association with a pharmaceutically acceptable carrier.

In accordance with the present invention there is provided a pharmaceutical composition for the preparation of a medicament for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells, which comprises

5 an INGAP peptide in an amount sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms; at least one immunosuppressive agent in an amount sufficient to protect said islet cells from immune destruction; and a pro-

10 survival factor in association with a pharmaceutically acceptable carrier.

In accordance with the present invention there is provided a pharmaceutical composition for the preparation of a medicament for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells, which comprises

15 an INGAP peptide in an amount sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms; and a pro-survival factor in association with a pharmaceutically acceptable carrier.

20 The immunosuppressive agent includes, without limitation, sirolimus, tacrolimus, or a combination thereof.

The term "pro-neogenesis factor" is intended to mean any compounds capable of islet regeneration including, without limitation, growth factors, GLP-1, exendin-4, and an INGAP peptide.

25 The preferred growth factors are selected from the group consisting of insulin, IGF-I, IGF-II, EGF, Gastrin and NGF.

The term "pro-survival factor" is intended to mean a factor including, without limitation, insulin, IGF-I, IGF-II, EGF and NGF.

30 The term "insulin-producing cells" is intended to mean pancreatic beta-cells.

The term "INGAP peptide" is intended to mean the fragment of native Islet Neogenesis Associated Protein (INGAP) protein which contains the biological activity of the full length molecule, including but not limited to, a biologically active fragment of:

35 Met Leu Pro Met Thr Leu Cys Arg Met Ser Trp Met Leu Leu Ser Cys  
 1 5 10 15

Leu Met Phe Leu Ser Trp Val Glu Gly Glu Glu Ser Gln Lys Lys Leu  
                     20                    25                    30  
 5 Pro Ser Ser Arg Ile Thr Cys Pro Gln Gly Ser Val Ala Tyr Gly Ser  
                     35                    40                    45  
 Tyr Cys Tyr Ser Leu Ile Leu Ile Pro Gln Thr Trp Ser Asn Ala Glu  
           50                    55                    60  
 10 Leu Ser Cys Gln Met His Phe Ser Gly His Leu Ala Phe Leu Leu Ser  
       65                    70                    75                    80  
 Thr Gly Glu Ile Thr Phe Val Ser Ser Leu Val Lys Asn Ser Leu Thr  
                     85                    90                    95  
 15 Ala Tyr Gln Tyr Ile Trp Ile Gly Leu His Asp Pro Ser His Gly Thr  
                     100                    105                    110  
 20 Leu Pro Asn Gly Ser Gly Trp Lys Trp Ser Ser Ser Asn Val Leu Thr  
           115                    120                    125  
 Phe Tyr Asn Trp Glu Arg Asn Pro Ser Ile Ala Ala Asp Arg Gly Tyr  
       130                    135                    140  
 25 Cys Ala Val Leu Ser Gln Lys Ser Gly Phe Gln Lys Trp Arg Asp Phe  
       145                    150                    155                    160  
 Asn Cys Glu Asn Glu Leu Pro Tyr Ile Cys Lys Phe Lys Val  
                     165                    170

30 (SEQ ID NO:1),  
 a fragment of 15 amino acids of the sequence SEQ ID NO: 1, more  
 precisely, such an INGAP peptide is of the following amino acid sequence:  
 Gly Leu His Asp Pro Ser His Gly Thr Leu Pro Asn Gly Ser Gly (SEQ ID  
 NO:2).

35 The term "islets of Langerhans" is intended to mean islet cells  
 and associated cells, such as duct cells, of any origin, such as human,  
 porcine, canine and murine, among others.

The term "neogenesis" is intended to mean the regeneration or  
*de novo* growth of cells.

40 Except as otherwise expressly defined herein, the abbreviations  
 used herein for designating the amino acids and the protective groups are  
 based on recommendations of the IUPAC-IUB Commission on Biochemical  
 Nomenclature (*Biochemistry*, 1972, 11:1726-1732).

### **BRIEF DESCRIPTION OF THE DRAWINGS**

Fig. 1 illustrates the increase in pancreatic insulin content and the reduction in the prevailing level of blood glucose resulting from the concurrent administration of INGAP peptide and sirolimus/tacrolimus and insulin in NOD mice.

Fig. 2 illustrates the survival of NOD mice treated with a combination of INGAP peptide, sirolimus/tacrolimus and insulin versus animals treated with sirolimus/tacrolimus alone or drug vehicle alone.

### **DETAILED DESCRIPTION OF THE INVENTION**

In accordance with the present invention, there is provided a method for the induction of *in vivo* islet cell neogenesis and new islet formation from cells derived from islet cell stem/progenitor cells in the adult pancreas, associated with the self-regulated expansion of such cells and the development of a mature glucose-sensing mechanism, leading to the reversal of an established diabetic state.

In accordance with one embodiment of the present invention, the technology is based on the understanding of autoimmune diabetes being a disease state characterized by a loss of an insulin-producing cell mass as a result of a pre-existing or ongoing autoimmune destruction of such cells, incorporating the following components that are necessary and sufficient for the successful reversal of a diabetic state by the induction of islet cell neogenesis and new islet formation:

1. a stimulus for the induction of islet cell neogenesis and new islet formation from pre-existing pancreatic stem/progenitor cells, provided by, but not limited to INGAP peptide;
2. provision of an immune tolerant environment to prevent ongoing or recurrent destruction of the newly regenerated cells, provided by, but not limited to, a combination of sirolimus/tacrolimus;
3. a pro-survival and anti-apoptosis factor, including but not limited to insulin.

The use of a pro-neogenesis factor is a critical part of the treatment, because without it, there is no stimulus to induce the transformation of putative stem/progenitor cells to new hormone-producing

islet cells. Alternatively, there may be such an endogenous stimulus but it may be ineffectual in terms of overcoming a much more effective ongoing cell destruction process. Hence it is the balance of neogenesis versus destruction that may be important.

5           Autoimmune diabetes, by definition, occurs through the autoimmune destruction of insulin-producing pancreatic beta-cells. In order to mitigate the ongoing or renewed destruction of such cells after the induction of islet cell neogenesis, the local immune environment must be altered to remove or diminish this autoimmune insult. Thus  
10 immunosuppressive agents, that include, but are not limited to a combination of sirolimus/tacrolimus, are required:

          Newly created beta-cells are known to be quite sensitive pro-death signals including, but not limited to high levels of circulating glucose. Thus pro-survival factors and in particular factors that can mitigate high  
15 levels of circulating glucose including, but not limited to insulin, are important to support and sustain cell survival.

          Evidence for the induction of islet cell neogenesis and new islet formation leading to the reversal of diabetes includes: (1) an increase in the expression of the transcription factor Pdx-1 in putative islet cell progenitor  
20 cells; (2) an increase in pancreatic insulin content; (3) an increase in beta-cell mass; (5) a decrease in the prevailing level of blood glucose; (6) an increase in survival.

          While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of  
25 further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential  
30 features hereinbefore set forth, and as follows in the scope of the appended claims.

**WHAT IS CLAIMED IS:**

1. A method to stimulate reversal of a diabetic state in a patient, which comprises *in vivo* inducing re-growth of new insulin-producing cells by administering a therapeutically effective amount of a pro-neogenesis factor to said patient, wherein formation of mature islets of Langerhans is indicative of a stimulated reversal of a diabetic state.
2. The method of claim 1, wherein said pro-neogenesis factor is selected from the group consisting of growth factors, GLP-1, exendin-4, and an INGAP peptide.
3. The method of claim 2, wherein said growth factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF, Gastrin and NGF.
4. The method of claim 1, wherein said insulin-producing cells are pancreatic beta-cells.
5. A method to prevent autoimmune destruction of new insulin-producing cells (pancreatic beta-cells) in a patient, which comprises administering to said patient a therapeutically effective amount of at least one immunosuppressive agent in combination with an INGAP peptide.
6. The method of claim 5, wherein said immunosuppressive agent is selected from the group consisting of sirolimus, tacrolimus, and a combination thereof.
7. A method to promote survival of the newly regenerated insulin-producing cells, which comprises administering a pro-neogenesis factor in a therapeutically effective amount to a patient.
8. The method of claim 7, wherein said pro-neogenesis factor is selected from the group consisting of growth factors, GLP-1, exendin-4, and an INGAP peptide.

9. The method of claim 8, wherein said growth factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF, Gastrin and NGF.

10. The method of claim 8, wherein said insulin-producing cells are pancreatic beta-cells.

11. An *in vivo* method for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells, which comprises the steps of:

- a) administering INGAP peptide to said patient in an amount sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms, whereby cells expand in number and develop a mature glucose-sensing mechanism in a regulated manner;
- b) concurrently administering to said patient at least one immunosuppressive agent in an amount sufficient to protect said islet cells from immune destruction; and
- c) concurrently administering a pro-survival factor to said patient during islet cell neogenesis and new islet formation.

12. The method of claim 11, wherein said islet hormone-producing cells are pancreatic beta-cells.

13. The method of claim 12, wherein said immunosuppressive agent is selected from the group consisting of sirolimus, tacrolimus, and a combination thereof.

14. The method of claim 11, wherein said pro-survival factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF and NGF.

15. An *in vivo* method for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells, which comprises the steps of:

- a) administering INGAP peptide to said patient in an amount sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms, whereby cells expand in number and develop a mature glucose-sensing mechanism in a regulated manner;
- b) concurrently administering a pro-survival factor to said patient during islet cell neogenesis and new islet formation.

16. The method of claim 15, wherein said islet hormone-producing cells are pancreatic beta-cells.

17. The method of claim 15, wherein said pro-survival factor is selected from the group consisting of insulin, IGF-I and IGF-II.

18. A pharmaceutical composition for the preparation of a medicament to stimulate reversal of a diabetic state in a patient by *in vivo* inducing re-growth of new insulin-producing cells, which comprises a therapeutically effective amount of a pro-neogenesis factor in association with a pharmaceutically acceptable carrier.

19. The pharmaceutical composition of claim 18, wherein said pro-neogenesis factor is selected from the group consisting of growth factors, GLP-1, exendin-4, and an INGAP peptide.

20. The pharmaceutical composition of claim 18, wherein said growth factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF, Gastrin and NGF.

21. A pharmaceutical composition for the preparation of a medicament to prevent autoimmune destruction of new insulin-producing cells in a patient, which comprises a therapeutically effective amount of at least one immunosuppressive agent and an INGAP peptide factor in association with a pharmaceutically acceptable carrier.



22. The pharmaceutical composition of claim 21, wherein said immunosuppressive agent is selected from the group consisting of sirolimus, tacrolimus, and a combination thereof.
23. A pharmaceutical composition for the preparation of a medicament to promote survival of the newly regenerated insulin-producing cells, which comprises a therapeutically effective amount of a pro-neogenesis factor in association with a pharmaceutically acceptable carrier.
24. The pharmaceutical composition of claim 23, wherein said pro-neogenesis factor is selected from the group consisting of growth factors, GLP-1, exendin-4, and an INGAP peptide.
25. The pharmaceutical composition of claim 24, wherein said growth factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF, Gastrin and NGF.
26. The pharmaceutical composition of claim 23, wherein said insulin-producing cells are pancreatic beta-cells.
27. A pharmaceutical composition for the preparation of a medicament for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells, which comprises an INGAP peptide in an amount sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms; at least one immunosuppressive agent in an amount sufficient to protect said islet cells from immune destruction; and a pro-survival factor in association with a pharmaceutically acceptable carrier.
28. The pharmaceutical composition of claim 27, wherein said islet hormone-producing cells are pancreatic beta-cells.

29. The pharmaceutical composition of claim 27, wherein said immunosuppressive agent is selected from the group consisting of sirolimus, tacrolimus, and a combination thereof.
30. The pharmaceutical composition of claim 29, wherein said pro-survival factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF and NGF.
31. A pharmaceutical composition for the preparation of a medicament for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells, which comprises an INGAP peptide in an amount sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms; and a pro-survival factor in association with a pharmaceutically acceptable carrier.
32. The pharmaceutical composition of claim 31, wherein said islet hormone-producing cells are pancreatic beta-cells.
33. The pharmaceutical composition of claim 31, wherein said pro-survival factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF, and NGF.
34. Use of a therapeutically effective amount of a pro-neogenesis factor to stimulate reversal of a diabetic state in a patient, wherein formation of mature islets of Langerhans is indicative of a stimulated reversal of a diabetic state.
35. The use of claim 34, wherein said pro-neogenesis factor is selected from the group consisting of growth factors, GLP-1, exendin-4, and an INGAP peptide.
36. The use of claim 35, wherein said growth factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF, Gastrin and NGF.

37. Use of a therapeutically effective amount of at least one immunosuppressive agent in combination with an INGAP peptide to prevent autoimmune destruction of new insulin-producing cells in a patient.
38. The use of claim 37, wherein said immunosuppressive agent is selected from the group consisting of sirolimus, tacrolimus, and a combination thereof.
39. Use of a therapeutically effective amount of a pro-neogenesis factor to promote survival of the newly regenerated insulin-producing cells.
40. The use of claim 39, wherein said pro-neogenesis factor is selected from the group consisting of growth factors, GLP-1, exendin-4, and an INGAP peptide.
41. The use of claim 40, wherein said growth factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF, Gastrin and NGF.
42. The use of claim 39, wherein said insulin-producing cells are pancreatic beta-cells.
43. Use of an INGAP peptide in an amount sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms; at least one immunosuppressive agent in an amount sufficient to protect said islet cells from immune destruction; and a pro-survival factor for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells in a patient.
44. The use of claim 43, wherein said islet hormone-producing cells are pancreatic beta-cells.

45. The use of claim 43, wherein said immunosuppressive agent is selected from the group consisting of sirolimus, tacrolimus, and a combination thereof.


46. The use of claim 43, wherein said pro-survival factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF and NGF.

47. Use of an INGAP peptide in an amount sufficient to stimulate transformation of putative islet cell stem/progenitor cells in adult pancreas into islet hormone-producing cells under normal endogenous homeostatic control mechanisms; and a pro-survival factor for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells in a patient.

48. The use of claim 47, wherein said islet hormone-producing cells are pancreatic beta-cells.

49. The use of claim 47, wherein said pro-survival factor is selected from the group consisting of insulin, IGF-I, IGF-II, EGF and NGF.

1/2



2/2




FIG. 2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
6 May 2004 (06.05.2004)

PCT

(10) International Publication Number  
**WO 2004/037277 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 38/10**,  
38/16, A61P 3/10

(21) International Application Number:  
PCT/CA2003/001635

(22) International Filing Date: 24 October 2003 (24.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/420,677 24 October 2002 (24.10.2002) US

(71) Applicant (for all designated States except US): **MCGILL UNIVERSITY** [CA/CA]; 845 Sherbrooke Street West, Montréal, Québec H3A 2T5 (CA).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **ROSENBERG, Lawrence** [CA/CA]; 6507 Fern Road, Côte St-Luc, Québec H4V 1E4 (CA).

(74) Agent: **OGILVY RENAULT**; Suite 1600, 1981 McGill College Avenue, Montreal, Québec H3A 2Y3 (CA).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:  
15 July 2004

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: USE OF INGP FOR REVERSING DIABETES

(57) Abstract: The present invention relates to a method to stimulate reversal of a diabetic state in a patient; a method to prevent autoimmune destruction of new insulin-producing cells (pancreatic beta-cells) in a patient; a method to promote survival of the newly regenerated insulin-producing cells (pancreatic beta-cells); and an in vivo method for the induction of islet cell neogenesis and new islet formation and the prevention of autoimmune destruction of said new cells.

WO 2004/037277 A3

## INTERNATIONAL SEARCH REPORT

 International Application No  
 PCT/03/01635

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K38/10 A61K38/16 A61P3/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, Sequence Search, BIOSIS, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 03/033808 A (PROCTER & GAMBLE) 24 April 2003 (2003-04-24)  * see claims 1 and 6, Seq ID No. 1-3 and examples *	1-4, 7-10, 18-20, 23-26, 34-36, 39-42
X	WO 02/070551 A (UNIV MCGILL ; MAYSINGER DUSICA (CA); ROSENBERG LAWRENCE (CA)) 12 September 2002 (2002-09-12) * see claims 1.-2 and 9, page 3 lines 38-41, page 6 lines 7-18, page 12 lines 11-15 *  ----- -/--	18-20, 23-26

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

17 May 2004

Date of mailing of the international search report

26/05/2004

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Merckling-Ruiz, V



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 03/01635

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96/26215 A (UNIV MCGILL ; EASTERN VIRGINIA MEDICAL SCHOO (US)) 29 August 1996 (1996-08-29)	1-4, 7-10, 18-20, 23-26, 34-36, 39-42
Y	* see claims 41-43 and Seq ID No 1-2 *	1-49
P,X	WO 03/057862 A (UNIV MCGILL ; ROSENBERG LAWRENCE (CA)) 17 July 2003 (2003-07-17)	1-4, 7-10, 18-20, 23-26, 34-36, 39-42
	* see claim 1 and paragraphs '20! and '21! *	
Y	WO 02/055152 A (WARATAH PHARMACEUTICALS INC) 18 July 2002 (2002-07-18) * see abstract, claim 1 and example 2 *	1-49
Y	RAFAELOFF RONIT ET AL: "Cloning and sequencing of the pancreatic islet neogenesis associated protein (INGAP) gene and its expression in islet neogenesis in hamsters" JOURNAL OF CLINICAL INVESTIGATION, NEW YORK, NY, US, vol. 99, no. 9, 1997, pages 2100-2109, XP002173530 ISSN: 0021-9738 * see abstract and page 2108 *	1-49

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/CA 03/01635

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-20, 23-26, 34-49  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 1-17 and 34-49 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 1-4, 7-10, 18-20, 23-26, 34-36, 39-42  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.1

Although claims 1-17 and 34-49 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

-----

## Continuation of Box I.2

Claims Nos.: 1-4,7-10,18-20,23-26,34-36,39-42

The application appears to be directed to compositions comprising an INGAP peptide, optionally in combination with other therapeutic agent(s), and their use in the treatment of diabetes (see pages 7-8 of the application). However, many claims are directed to the use of a "proneogenesis factor", which is defined in a much broader manner (see dependent claims 2 and 3). These embodiments are not supported at all by the application, not even in a theoretical manner. Therefore, claims 1-4,7-10,18-20,23-26,34-36,39-42 partially lack disclosure in the sense of Art. 5 PCT. These claims were only searched insofar as the "proneogenesis factor" is INGAP.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.5), should the problems which led to the Article 17(2) declaration be overcome.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 03/01635

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03033808	A	24-04-2003	WO 03033808 A2	24-04-2003
WO 02070551	A	12-09-2002	WO 02070551 A2	12-09-2002
WO 9626215	A	29-08-1996	US 5834590 A	10-11-1998
			AU 708499 B2	05-08-1999
			AU 4914996 A	11-09-1996
			CA 2213610 A1	29-08-1996
			EP 0815129 A1	07-01-1998
			JP 11500907 T	26-01-1999
			WO 9626215 A1	29-08-1996
			US 5840531 A	24-11-1998
WO 03057862	A	17-07-2003	WO 03057862 A2	17-07-2003
			US 2003203850 A1	30-10-2003
WO 02055152	A	18-07-2002	CA 2434330 A1	18-07-2002
			CN 1486204 T	31-03-2004
			EP 1351742 A2	15-10-2003
			WO 02055152 A2	18-07-2002
			US 2002098178 A1	25-07-2002

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: \_\_\_\_\_**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**